

# Summary

Production Name	Oct-2 Rabbit Polyclonal Antibody	
Description	Rabbit Polyclonal Antibody	
Host	Rabbit	
Application	IF,WB,ELISA	
Reactivity	Human, Mouse, Monkey	

## Performance

Conjugation	Unconjugated	
Modification	Unmodified	
lsotype	lgG	
Clonality	Polyclonal	
Form	Liquid	
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw	
	cycles.	
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.	
Purification	Affinity purification	

### Immunogen

Gene Name	POU2F2		
	POU2F2; OCT2; OTF2; POU domain; class 2, transcription factor 2; Lymphoid-restricted		
Alternative Names	immunoglobulin octamer-binding protein NF-A2; Octamer-binding protein 2; Oct-2;		
	Octamer-binding transcription factor 2; OTF-2		
Gene ID	5452.0		
SwissProt ID	P09086.The antiserum was produced against synthesized peptide derived from human		
	OCT2. AA range:1-50		

# Application

Dilution Ratio	WB 1:500 - 1:2000.	IF 1:200 - 1:1000. ELISA	A: 1:20000. Not yet tested in other
Dilution Ratio	1		
	applications.		

## Product Name: Oct-2 Rabbit Polyclonal Antibody Catalog #: APRab15102




Molecular Weight 52kD

## Background

The protein encoded by this gene is a homeobox-containing transcription factor of the POU domain family. The encoded protein binds the octamer sequence 5'-ATTTGCAT-3', a common transcription factor binding site in immunoglobulin gene promoters. Several transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2011],function:Transcription factor that specifically binds to the octamer motif (5'-ATTTGCAT-3'). Regulates transcription in a number of tissues in addition to activating immunoglobulin gene expression. Modulates transcription transactivation by NR3C1, AR and PGR. Isoform 5 activates the U2 small nuclear RNA (snRNA) promoter.,similarity:Belongs to the POU transcription factor family. Class-2 subfamily.,similarity:Contains 1 homeobox DNA-binding domain.,similarity:Contains 1 POU-specific domain.,subunit:Interacts with NR3C1, AR and PGR.,tissue specificity:Isoform 3 is B-cell specific. Isoform 5 is expressed in B-cells and the immunoglobulin-expressing T-cell line Molt-4, but not in the T-cell line BW 5147.,


## **Research Area**

#### Image Data




Immunofluorescence analysis of COS7 cells, using OCT2 Antibody. The picture on the right is blocked with the synthesized peptide.





Western blot analysis of lysates from COS7 cells, using OCT2 Antibody. The lane on the right is blocked with the synthesized



Western Blot analysis of various cells using Oct-2 Polyclonal Antibody

Note

For research use only.