

## Summary

Olfactory receptor 10J3 Rabbit Polyclonal Antibody	
Rabbit Polyclonal Antibody	
Rabbit	
WB,ELISA	
Human, Rat, Mouse	
Rabbit Polyclonal Antibody Rabbit WB,ELISA	

#### Performance

Conjugation	Unconjugated	
Modification	Unmodified	
lsotype	lgG	
Clonality	Polyclonal	
Form	Liquid	
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw	
Storage	cycles.	
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.	
Purification	Affinity purification	

#### Immunogen

Gene Name	OR10J3	
Alternative Names	OR10J3; OR10J3P; Olfactory receptor 10J3	
Gene ID	441911.0	
SwissProt ID	Q5JRS4.The antiserum was produced against synthesized peptide derived from human	
	OR10J3. AA range:237-286	

# Application

Dilution Ratio	WB 1:500 - 1:2000. ELISA: 1:20000
Molecular Weight	37kD




#### Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,


#### **Research Area**

Olfactory transduction;

### Image Data



Western blot analysis of lysates from MCF-7 cells, using OR10J3 Antibody. The lane on the right is blocked with the



Western blot analysis of the lysates from HUVECcells using OR10J3 antibody.



#### Note

For research use only.