

# Summary

Production Name	Olfactory receptor 52D1 Rabbit Polyclonal Antibody			
Description	Rabbit Polyclonal Antibody			
Host	Rabbit			
Application	IF,WB,			
Reactivity	Human, Monkey			

#### Performance

Conjugation	Unconjugated		
Modification	Unmodified		
lsotype	IgG		
Clonality	Polyclonal		
Form	Liquid		
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw		
	cycles.		
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.		
Purification	Affinity purification		

#### Immunogen

Gene Name	OR52D1
Alternative Names	OR52D1; Olfactory receptor 52D1; Odorant receptor HOR5'beta14; Olfactory receptor
	OR11-43
Gene ID	390066.0
SwissProt ID	Q9H346.The antiserum was produced against synthesized peptide derived from human
	OR52D1. AA range:269-318

# Application

Dilution Ratio	WB 1:500 - 1:2000.	. IF 1:200 - 1	:1000. ELISA: 1:5000.	Not yet tested in other
	applications.			

## Product Name: Olfactory receptor 52D1 Rabbit Polyclonal Antibody Catalog #: APRab15260



Molecular Weight 35kD


## Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,


### **Research Area**

Olfactory transduction;

## Image Data




Immunofluorescence analysis of MCF7 cells, using OR52D1 Antibody. The picture on the right is blocked with the




Western blot analysis of lysates from COLO and COS7 cells, using OR52D1 Antibody. The lane on the right is blocked with





Western blot analysis of the lysates from HepG2 cells using OR52D1 antibody.



Western Blot analysis of various cells using Olfactory receptor 52D1 Polyclonal Antibody

#### Note

For research use only.