

Catalog #: APRab15319



Summary

Olfactory receptor 7C2 Rabbit Polyclonal Antibody **Production Name**

Description Rabbit Polyclonal Antibody

Rabbit Host **Application** WB.ELISA

Reactivity Human, Rat, Mouse

Performance

Conjugation Unconjugated Modification Unmodified

IgG Isotype

Clonality Polyclonal **Form** Liquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw Storage

cycles.

Buffer Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

Purification Affinity purification

Immunogen

Gene Name OR7C2

OR7C2; OR7C3; Olfactory receptor 7C2; Olfactory receptor 19-18; OR19-18; Olfactory **Alternative Names**

receptor 7C3; Olfactory receptor OR19-22

Gene ID 26658.0

O60412. The antiserum was produced against synthesized peptide derived from human SwissProt ID

OR7C2. AA range:198-247

Application

Dilution Ratio WB 1:500 - 1:2000. ELISA: 1:10000. Not yet tested in other applications.

Molecular Weight 35kD

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Antibody

Catalog #: APRab15319




Background

olfactory receptor family 7 subfamily C member 2(OR7C2) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008], function: Odorant receptor ., similarity: Belongs to the G-protein coupled receptor 1 family.,


Research Area

Olfactory transduction;

Image Data



Western blot analysis of lysates from HUVEC cells, using OR7C2 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HeLa cells using OR7C2 antibody.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838



HuvEc (kD)
11785483426-

19-

Western Blot analysis of various cells using Olfactory receptor 7C2 Polyclonal Antibody

Note

For research use only.