

Summary

Olfactory receptor 8K3 Rabbit Polyclonal Antibody
Rabbit Polyclonal Antibody
Rabbit
IF,WB,ELISA
Human, Rat, Mouse

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storege	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
Storage	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	OR8K3
Alternative Names	OR8K3; Olfactory receptor 8K3; Olfactory receptor OR11-181
Gene ID	219473.0
SwissProt ID	Q8NH51.The antiserum was produced against synthesized peptide derived from
	human OR8K3. AA range:236-285

Application

Dilution Ratio	WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:10000. Not yet tested in other
	applications.
Molecular Weight	33kD

Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms. This olfactory receptor gene is a segregating pseudogene, where some individuals have an allele that encodes a functional olfactory receptor, while other individuals have an allele encoding afunction:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,


Research Area

Olfactory transduction;

Image Data




Immunofluorescence analysis of HUVEC cells, using OR8K3 Antibody. The picture on the right is blocked with the







Western blot analysis of lysates from LOVO cells, using OR8K3 Antibody. The lane on the right is blocked with the



Western Blot analysis of LOVO cells using Olfactory receptor 8K3 Polyclonal Antibody

Note

For research use only.