

Summary

| Production Name | p70 S6 kinase α Rabbit Polyclonal Antibody | | |
|-----------------|---|--|--|
| Description | Rabbit Polyclonal Antibody | | |
| Host | Rabbit | | |
| Application | IF,IHC,WB, | | |
| Reactivity | Human,Mouse,Rat,Pig | | |

Performance

| Conjugation | Unconjugated | | |
|--------------|--|--|--|
| Modification | Unmodified | | |
| lsotype | lgG | | |
| Clonality | Polyclonal | | |
| Form | Liquid | | |
| Storage | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw | | |
| | cycles. | | |
| Buffer | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N. | | |
| Purification | Affinity purification | | |

Immunogen

| Gene Name | RPS6KB1 | |
|-------------------|---|--|
| Alternative Names | RPS6KB1; STK14A; Ribosomal protein S6 kinase beta-1; S6K-beta-1; S6K1; 70 kDa | |
| | ribosomal protein S6 kinase 1; P70S6K1; p70-S6K 1; Ribosomal protein S6 kinase I; | |
| | Serine/threonine-protein kinase 14A; p70 ribosomal S6 kinase alpha; p70 S6 kinas | |
| Gene ID | 6198.0 | |
| SwissProt ID | P23443.The antiserum was produced against synthesized peptide derived from human | |
| | p70 S6 Kinase. AA range:337-386 | |

Application

| Dilution Ratio | WB 1:500 - 1:2000 | IHC 1:100 - 1:300. IF 1:200 | - 1:1000. ELISA: 1:20000. Not yet tested |
|----------------|-------------------|-----------------------------|--|
|----------------|-------------------|-----------------------------|--|



in other applications.

Molecular Weight

60kD


Background

ribosomal protein S6 kinase B1(RPS6KB1) Homo sapiens This gene encodes a member of the ribosomal S6 kinase family of serine/threonine kinases. The encoded protein responds to mTOR (mammalian target of rapamycin) signaling to promote protein synthesis, cell growth, and cell proliferation. Activity of this gene has been associated with human cancer. Alternatively spliced transcript variants have been observed. The use of alternative translation start sites results in isoforms with longer or shorter N-termini which may differ in their subcellular localizations. There are two pseudogenes for this gene on chromosome 17. [provided by RefSeq, Jan 2013],catalytic activity:ATP + a protein = ADP + a phosphoprotein.,enzyme regulation:Activation by serine/threonine phosphorylation and protein kinase C, inactivated by type 2A phosphatase, function:Phosphorylates specifically ribosomal protein S6 in response to insulin or several classes of mitogens.,similarity:Belongs to the protein kinase superfamily, similarity:Belongs to the protein kinase superfamily, similarity:Contains 1 AGC-kinase C-terminal domain.,similarity:Contains 1 protein kinase domain.,subunit:Interacts with PPP1R9A/neurabin-1,tissue specificity:Widely expressed.,

Research Area


Regulates Angiogenesis; Insulin Receptor; ErbB/HER; mTOR; B Cell Receptor; PI3K/Akt; PI3K/Akt; AMPK

Image Data




Immunofluorescence analysis of NIH/3T3 cells, using p70 S6 Kinase Antibody. The picture on the right is blocked with the synthesized peptide.






Immunohistochemistry analysis of paraffin-embedded human brain tissue, using p70 S6 Kinase Antibody. The picture on the



Western blot analysis of lysates from HeLa/HepG2, using p70 S6 Kinase Antibody. The lane on the right is blocked with the



Western Blot analysis of various cells using p70 S6 kinase α Polyclonal Antibody

Note

For research use only.