

# Summary

| Production Name | PIAS 3 Rabbit Polyclonal Antibody |
|-----------------|-----------------------------------|
| Description     | Rabbit Polyclonal Antibody        |
| Host            | Rabbit                            |
| Application     | IHC,WB,                           |
| Reactivity      | Human, Mouse, Rat                 |

#### Performance

| Conjugation  | Unconjugated   |
|--------------|--|
| Modification | Unmodified   |
| lsotype      | IgG  |
| Clonality    | Polyclonal   |
| Form         | Liquid   |
| Storage      | Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw   |
|              | cycles.  |
| Buffer       | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N. |
| Purification | Affinity purification  |

#### Immunogen

| Gene Name         | PIAS3  |
|-------------------|--|
| Alternative Names | PIAS3; E3 SUMO-protein ligase PIAS3; Protein inhibitor of activated STAT protein 3 |
| Gene ID           | 10401.0  |
| SwissProt ID      | Q9Y6X2. The antiserum was produced against synthesized peptide derived from human  |
|                   | PIAS3. AA range:10-59  |

# Application

| Dilution Ratio   | WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:10000 |
|------------------|--|
| Molecular Weight | 68kD   |

## Background

## Product Name: PIAS 3 Rabbit Polyclonal Antibody Catalog #: APRab16120



This gene encodes a member of the PIAS [protein inhibitor of activated STAT (signal transducer and activator of transcription)] family of transcriptional modulators. The protein functions as a SUMO (small ubiquitin-like modifier)-E3 ligase which catalyzes the covalent attachment of a SUMO protein to specific target substrates. It directly binds to several transcription factors and either blocks or enhances their activity. Alternatively spliced transcript variants of this gene have been identified, but the full-length nature of some of these variants has not been determined. [provided by RefSeq, Jul 2008],domain:The LXXLL motif is a transcriptional coregulator signature.,function:Functions as an E3-type small ubiquitin-like modifier (SUMO) ligase, stabilizing the interaction between UBE2I and the substrate, and as a SUMO-tethering factor. Plays a crucial role as a transcriptional coregulation in various cellular pathways, including the STAT pathway and the steroid hormone signaling pathway. The effects of this transcriptional coregulation, transactivation or silencing, may vary depending upon the biological context.,induction:By dihydrotestosterone in prostate cancer cells.,pathway:Protein modification; protein sumoylation.,PTM:Sumoylated.,similarity:Belongs to the PIAS family.,similarity:Contains 1 SAP domain.,similarity:Contains 1 SP-RING-type zinc finger.,subunit:Binds SUMO1 and UBE2I. Interacts with AR, GFI1, HMGA2, IRF1, MITF, NCOA2, as well as with STAT3, after treatment with IL6, CNTF or OSM and with STAT5, after PRL stimulation (By similarity). Interacts with PLAG1.,tissue specificity:Widely expressed.,

## **Research Area**

Ubiquitin mediated proteolysis;Jak\_STAT;Pathways in cancer;Small cell lung cancer;

# Image Data



Immunohistochemistry analysis of paraffin-embedded human lung carcinoma tissue, using PIAS3 Antibody. The picture on the right is blocked with the synthesized peptide.





Western blot analysis of lysates from 293 cells, treated with UV 5 ', using PIAS3 Antibody. The lane on the right is blocked



Western Blot analysis of various cells using PIAS 3 Polyclonal Antibody diluted at 1: 2000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



Western Blot analysis of 293 cells using PIAS 3 Polyclonal Antibody diluted at 1: 2000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).

#### Note

For research use only.