

Product Name: PRKX Rabbit Polyclonal Antibody
Catalog #: AP Rab16500



Summary

Production Name	PRKX Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IHC, WB, ELISA
Reactivity	Human, Mouse, Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	PRKX PRKX; PKX1; cAMP-dependent protein kinase catalytic subunit PRKX; PrKX; Protein kinase X; Protein kinase X-linked; Serine/threonine-protein kinase PRKX; Protein kinase PKX1
Alternative Names	
Gene ID	5613.0
SwissProt ID	P51817. The antiserum was produced against synthesized peptide derived from human PRKX. AA range: 251-300

Application

Dilution Ratio	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000..
Molecular Weight	41kD


Background

This gene encodes a serine threonine protein kinase that has similarity to the catalytic subunit of cyclic AMP dependent protein kinases. The encoded protein is developmentally regulated and may be involved in renal epithelial morphogenesis. This protein may also be involved in macrophage and granulocyte maturation. Abnormal recombination between this gene and a related pseudogene on chromosome Y is a frequent cause of sex reversal disorder in XX males and XY females. Pseudogenes of this gene are found on chromosomes X, 15 and Y. [provided by RefSeq, Feb 2010], catalytic activity: ATP + a protein = ADP + a phosphoprotein., disease: A chromosomal aberration involving PRKX is a cause of sex reversal disorder. Translocation t(X;Y)(p22;p11) with PRKY. Chromosomal translocations proximal to PRKY account for about 30% of the cases of sex reversal disorder in XX males and XY females., similarity: Belongs to the protein kinase superfamily. AGC Ser/Thr protein kinase family. cAMP subfamily., similarity: Contains 1 AGC-kinase C-terminal domain., similarity: Contains 1 protein kinase domain., tissue specificity: High levels in adult and fetal brain, kidney and lung; low levels in adult placenta, heart, liver, skeletal muscle, pancreas and fetal liver.,

Research Area


MAPK_ERK_Growth; MAPK_G_Protein; Calcium; Chemokine; Oocyte meiosis; Apoptosis_Inhibition; Apoptosis_Mitochondrial; Apoptosis_Overview; Vascular smooth muscle contraction; WNT; WNT-T CELL Hedgehog; Gap junction; Long-term potentiation; Olfactory transduction; Taste transduction; Insulin_Receptor; GnRH; Progesterone-mediated oocyte maturation; Melanogenesis; Prion diseases; Vibrio cholerae infection; Dilated cardiomyopathy;

Image Data




Immunohistochemistry analysis of paraffin-embedded human breast carcinoma tissue, using PRKX Antibody. The picture on the right is blocked with the synthesized peptide.

Product Name: PRKX Rabbit Polyclonal Antibody
Catalog #: AP Rab16500



Western blot analysis of lysates from MCF-7 cells, using PRKX Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HepG2 cells using PRKX antibody.

Note

For research use only.