

Product Name: RGS1 Rabbit Polyclonal Antibody
Catalog #: APRab17088



Summary

Production Name	RGS1 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IHC, WB,
Reactivity	Human, Mouse, Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	RGS1
Alternative Names	RGS1; 1R20; BL34; IER1; Regulator of G-protein signaling 1; RGS1; B-cell activation protein BL34; Early response protein 1R20
Gene ID	5996.0
SwissProt ID	Q08116. The antiserum was produced against synthesized peptide derived from human RGS1. AA range: 118-167

Application


Dilution Ratio	WB 1:500 - 1:2000. IHC 1:100 - 1:300. ELISA: 1:20000..
Molecular Weight	22kD

Background


This gene encodes a member of the regulator of G-protein signalling family. This protein is located on the cytosolic side of the plasma membrane and contains a conserved, 120 amino acid motif called the RGS domain. The protein attenuates the signalling activity of G-proteins by binding to activated, GTP-bound G alpha subunits and acting as a GTPase activating protein (GAP), increasing the rate of conversion of the GTP to GDP. This hydrolysis allows the G alpha subunits to bind G beta/gamma subunit heterodimers, forming inactive G-protein heterotrimers, thereby terminating the signal. [provided by RefSeq, Jul 2008],function:Inhibits signal transduction by increasing the GTPase activity of G protein alpha subunits thereby driving them into their inactive GDP-bound form. This protein may be involved in the regulation of B-cell activation and proliferation.,induction:In response to several B-cell activation signals,PTM:Could be phosphorylated. Might be functionally regulated by protein kinase(s),similarity:Contains 1 RGS domain.,tissue specificity:B-cell specific. Expression is relatively low in B-cells and chronic lymphocytic leukemia B-cells; however, in other types of malignant B-cell such as non-Hodgkin's lymphoma and hairy cell leukemia, expression is constitutively high.,

Research Area

Image Data




Immunohistochemistry analysis of paraffin-embedded human colon carcinoma tissue, using RGS1 Antibody. The picture on the right is blocked with the synthesized peptide.




Product Name: RGS1 Rabbit Polyclonal Antibody
Catalog #: APRab17088



Western blot analysis of lysates from mouse muscle cells, using RGS1 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from 293 cells using RGS1 antibody.



Western Blot analysis of various cells using RGS1 Polyclonal Antibody

Note

For research use only.