

Summary

Production Name	RNase III Drosha Rabbit Polyclonal Antibody	
Description	Rabbit Polyclonal Antibody	
Host	Rabbit	
Application	WB,IHC,ELISA	
Reactivity	Human, Mouse	

Performance

Conjugation	Unconjugated
Modification	Unmodified
lsotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw
	cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	DROSHA
Alternative Names	DROSHA; RN3; RNASE3L; RNASEN; Ribonuclease 3; Protein Drosha; Ribonuclease III;
	RNase III; p241
Gene ID	29102.0
SwissProt ID	Q9NRR4.The antiserum was produced against synthesized peptide derived from
	human RNase III Drosha. AA range:774-823

Application

Dilution Ratio	WB 1:500 - 1:2000. IHC: 1:100-1:300. ELISA: 1:10000
Molecular Weight	160kD



Background

drosha ribonuclease III(DROSHA) Homo sapiens This gene encodes a ribonuclease (RNase) III double-stranded RNAspecific ribonuclease and subunit of the microprocessor protein complex, which catalyzes the initial processing step of microRNA (miRNA) synthesis. The encoded protein cleaves the stem loop structure from the primary microRNA (primiRNA) in the nucleus, yielding the precursor miRNA (pre-miRNA), which is then exported to the cytoplasm for further processing. In a human cell line lacking a functional copy of this gene, canonical miRNA synthesis is reduced. Somatic mutations in this gene have been observed in human patients with kidney cancer. [provided by RefSeq, Sep 2016],catalytic activity:Endonucleolytic cleavage to 5'-phosphomonoester.,cofactor:Magnesium or manganese.,function:Executes the initial step of microRNA (miRNA) processing in the nucleus, that is cleavage of pri-miRNA to release pre-miRNA. Involved in prerRNA processing. Cleaves double-strand RNA and does not cleave single-strand RNA,online information:The dark side of RNA -Issue 87 of October 2007,similarity:Contains 1 DRBM (double-stranded RNA-binding) domain.,similarity:Contains 2 RNase III domains.,subcellular location:A fraction is translocated to the nucleolus during the S phase of the cell cycle.,subunit:Interacts with Sp1,tissue specificity:Ubiquitous.,

Research Area

Image Data



Western blot analysis of lysates from brain tissue, using RNase III Drosha antibody.



SH-SY5Y K562 HELA		
138: 100 70	RNase III Drosha	
55		
40		
35		
25		
15		

Western blot analysis of SH-SY5Y K562 HELA using RNase III Drosha antibody. Antibody was diluted at 1:1000. Secondary antibody was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-



Immunohistochemical analysis of paraffin-embedded human-brain, antibody was diluted at 1:200



Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:200

Product Name: RNase III Drosha Rabbit Polyclonal Antibody Catalog #: APRab17277





Immunohistochemical analysis of paraffin-embedded human-kidney, antibody was diluted at 1:200

Note For research use only.