

**Product Name: SEC61B Rabbit Polyclonal Antibody**  
**Catalog #: APRab17695**



## Summary

<b>Production Name</b>	SEC61B Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,ELISA
<b>Reactivity</b>	Human,Mouse,Rat

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	SEC61B
<b>Alternative Names</b>	Protein transport protein Sec61 subunit beta
<b>Gene ID</b>	10952.0
<b>SwissProt ID</b>	P60468.Synthesized peptide derived from human SEC61B AA range: 30-110

## Application

<b>Dilution Ratio</b>	IHC 1:50-200 ELISA(peptide)1:5000-20000
<b>Molecular Weight</b>	

## Background

The Sec61 complex is the central component of the protein translocation apparatus of the endoplasmic reticulum (ER) membrane. Oligomers of the Sec61 complex form a transmembrane channel where proteins are translocated across and

**Product Name: SEC61B Rabbit Polyclonal Antibody**  
**Catalog #: APRab17695**


---



integrated into the ER membrane. This complex consists of three membrane proteins- alpha, beta, and gamma. This gene encodes the beta-subunit protein. The Sec61 subunits are also observed in the post-ER compartment, suggesting that these proteins can escape the ER and recycle back. There is evidence for multiple polyadenylated sites for this transcript. [provided by RefSeq, Jul 2008],function:Necessary for protein translocation in the endoplasmic reticulum.,similarity:Belongs to the SEC61-beta family.,subunit:Heterotrimeric complex composed of SEC61-alpha, SEC61-beta and SEC61-gamma. Part of a complex composed of SEC61, SEC62 and SEC63. Interacts with SEC62.,

## Research Area

## Image Data



Immunohistochemical analysis of paraffin-embedded human Colon cancer. 1, Antibody was diluted at 1:200 (4° overnight) .  
2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200 (room temperature, 45min) .

## Note

For research use only.