

Product Name: SF-1 Rabbit Polyclonal Antibody
Catalog #: APRab17780



Summary

Production Name	SF-1 Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	NR5A1 NR5A1; AD4BP; FTZF1; SF1; Steroidogenic factor 1; SF-1; STF-1; Adrenal 4-binding protein; Fushi tarazu factor homolog 1; Nuclear receptor subfamily 5 group A member 1; Steroid hormone receptor Ad4BP
Alternative Names	
Gene ID	2516.0
SwissProt ID	Q13285.The antiserum was produced against synthesized peptide derived from human NR5A1. AA range:169-218

Application


Dilution Ratio	WB 1:500-2000 ELISA 2000-20000
Molecular Weight	50kD

Background

The protein encoded by this gene is a transcriptional activator involved in sex determination. The encoded protein binds DNA as a monomer. Defects in this gene are a cause of XY sex reversal with or without adrenal failure as well as adrenocortical insufficiency without ovarian defect. [provided by RefSeq, Jul 2008],disease:Defects in NR5A1 are a cause of adrenocortical insufficiency without ovarian defect [MIM:184757]. The disease is characterized by severe 'slackness,' muscular hypotonia. There is decreased sodium, increased potassium and elevated ACTH.,disease:Defects in NR5A1 are a cause of XY sex reversal with or without adrenal failure [MIM:184757]. This disease is characterized by normal female external genitalia and retention of the uterus.,function:Transcriptional activator. Seems to be essential for sexual differentiation and formation of the primary steroidogenic tissues. Binds to the Ad4 site found in the promoter region of steroidogenic P-450 genes such as CYP11A, CYP11B and CYP21B. Also regulates the Muellerian inhibiting substance (AMH) gene as well as the AHCH and STAR genes. 5'-YCAAGGYC-3' and 5'-RRAGGTCA-3' are the consensus sequences for the recognition by NR5A1/FTZF1. The SFPQ-NONO-NR5A1/SF-1 complex binds to the CYP17 promoter and regulates basal and cAMP-dependent transcriptional activity. Binds phosphatidylcholine (By similarity). Binds phospholipids with a phosphatidylinositol (PI) headgroup, in particular PI(3,4)P2 and PI(3,4,5)P3.,PTM:Acetylation stimulates the transcriptional activity.,similarity:Belongs to the nuclear hormone receptor family.,similarity:Belongs to the nuclear hormone receptor family. NR5 subfamily.,similarity:Contains 1 nuclear receptor DNA-binding domain.,subunit:Binds DNA as a monomer. Interacts with NR0B2 and PPARGC1A (By similarity). Part of a complex consisting of SFPQ, NONO and NR5A1/SF-1. Interacts with NCOA2.,

Research Area

Image Data



Western blot analysis of lysates from rat lung, using STF-1 Antibody. The lane on the right is blocked with the synthesized peptide.

Product Name: SF-1 Rabbit Polyclonal Antibody
Catalog #: APRab17780



Note

For research use only.