

**Product Name: SNAT2 Rabbit Polyclonal Antibody**  
**Catalog #: APRab18051**

---



## Summary

<b>Production Name</b>	SNAT2 Rabbit Polyclonal Antibody
<b>Description</b>	Rabbit Polyclonal Antibody
<b>Host</b>	Rabbit
<b>Application</b>	WB,ELISA
<b>Reactivity</b>	Human,Rat,Mouse

## Performance

<b>Conjugation</b>	Unconjugated
<b>Modification</b>	Unmodified
<b>Isotype</b>	IgG
<b>Clonality</b>	Polyclonal
<b>Form</b>	Liquid
<b>Storage</b>	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
<b>Buffer</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
<b>Purification</b>	Affinity purification

## Immunogen

<b>Gene Name</b>	SLC38A2
<b>Alternative Names</b>	SLC38A2; ATA2; KIAA1382; SAT2; SNAT2; Sodium-coupled neutral amino acid transporter 2; Amino acid transporter A2; Protein 40-9-1; Solute carrier family 38 member 2; System A amino acid transporter 2; System A transporter 1; System N amino a
<b>Gene ID</b>	54407.0
<b>SwissProt ID</b>	Q96QD8.The antiserum was produced against synthesized peptide derived from human SLC38A2. AA range:151-200

## Application

<b>Dilution Ratio</b>	WB 1:500-2000 ELISA 2000-20000
<b>Molecular Weight</b>	50kD

---


## Background

enzyme regulation: Inhibited by N-methyl-D-glucamine and probably choline., function: Functions as a sodium-dependent amino acid transporter. Mediates the saturable, pH-sensitive and electrogenic cotransport of neutral amino acids and sodium ions with a stoichiometry of 1:1. May function in the transport of amino acids at the blood-brain barrier and in the supply of maternal nutrients to the fetus through the placenta., induction: Up-regulated upon hypertonic conditions and amino acid deprivation., miscellaneous: Depletion of SLC38A2 by siRNA prevents the recovery of cells from hypertonic stress., PTM: Polyubiquitination by NEDD4L regulates the degradation and the activity of SLC38A2., similarity: Belongs to the amino acid/polyamine transporter 2 family., subcellular location: Insulin promotes recruitment to the plasma membrane from a pool localized in the trans-Golgi network or endosomes (By similarity). Enriched in the somatodendritic compartment of neurons, it is also detected at the axonal shaft but excluded from the nerve terminal., tissue specificity: Ubiquitously expressed. Widely expressed in the central nervous system with higher concentrations in caudal regions. Expressed by glutamatergic and GABAergic neurons together with astrocytes and other non-neuronal cells in the cerebral cortex (at protein level)., enzyme regulation: Inhibited by N-methyl-D-glucamine and probably choline., function: Functions as a sodium-dependent amino acid transporter. Mediates the saturable, pH-sensitive and electrogenic cotransport of neutral amino acids and sodium ions with a stoichiometry of 1:1. May function in the transport of amino acids at the blood-brain barrier and in the supply of maternal nutrients to the fetus through the placenta., induction: Up-regulated upon hypertonic conditions and amino acid deprivation., miscellaneous: Depletion of SLC38A2 by siRNA prevents the recovery of cells from hypertonic stress., PTM: Polyubiquitination by NEDD4L regulates the degradation and the activity of SLC38A2., similarity: Belongs to the amino acid/polyamine transporter 2 family., subcellular location: Insulin promotes recruitment to the plasma membrane from a pool localized in the trans-Golgi network or endosomes (By similarity). Enriched in the somatodendritic compartment of neurons, it is also detected at the axonal shaft but excluded from the nerve terminal., tissue specificity: Ubiquitously expressed. Widely expressed in the central nervous system with higher concentrations in caudal regions. Expressed by glutamatergic and GABAergic neurons together with astrocytes and other non-neuronal cells in the cerebral cortex (at protein level).


## Research Area

## Image Data

**Product Name: SNAT2 Rabbit Polyclonal Antibody**  
**Catalog #: APRab18051**



Western blot analysis of lysates from HeLa cells, using SLC38A2 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from HeLa cells using SLC38A2 antibody.

## Note

For research use only.