

Product Name: Synuclein- β Rabbit Polyclonal Antibody
Catalog #: APRab18518



Summary

Production Name	Synuclein- β Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	IF,IHC,WB,
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	SNCB
Alternative Names	SNCB; Beta-synuclein
Gene ID	6620.0
SwissProt ID	Q16143.The antiserum was produced against synthesized peptide derived from human Synuclein beta. AA range:85-134

Application


Dilution Ratio	WB 1:500 - 1:2000 IHC 1:100 - 1:300. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other applications.
Molecular Weight	14kD

Background

This gene encodes a member of a small family of proteins that inhibit phospholipase D2 and may function in neuronal plasticity. The encoded protein is abundant in lesions of patients with Alzheimer disease. A mutation in this gene was found in individuals with dementia with Lewy bodies. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2015], disease: Brain iron accumulation type 1 (NBIA1, also called Hallervorden-Spatz syndrome), a rare neuroaxonal dystrophy, is histologically characterized by axonal spheroids, iron deposition, Lewy body (LB)-like intraneuronal inclusions, glial inclusions and neurofibrillary tangles. SNCB is found in spheroids but not in inclusions., function: Non-amyloid component of senile plaques found in Alzheimer disease. Could act as a regulator of SNCA aggregation process. Protects neurons from staurosporine and 6-hydroxy dopamine (6OHDA)-stimulated caspase activation in a TP53/p53-dependent manner. Contributes to restore the SNCA anti-apoptotic function abolished by 6OHDA. Not found in the Lewy bodies associated with Parkinson disease., PTM: Phosphorylated. Phosphorylation by G-protein coupled receptor kinases (GRK) is more efficient than phosphorylation by CK1, CK2 and CaM-kinase II., similarity: Belongs to the synuclein family., tissue specificity: Expressed predominantly in brain; concentrated in presynaptic nerve terminals.,


Research Area

Image Data




Immunofluorescence analysis of A549 cells, using Synuclein beta Antibody. The picture on the right is blocked with the synthesized peptide.

Product Name: Synuclein- β Rabbit Polyclonal Antibody
Catalog #: APRab18518



Immunohistochemistry analysis of paraffin-embedded human brain tissue, using Synuclein beta Antibody. The picture on the right is blocked with the synthesized peptide.



Western blot analysis of lysates from HeLa cells, using Synuclein beta Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using Synuclein- β Polyclonal Antibody

Note

For research use only.