# **Product Name: T2R13 Rabbit Polyclonal Antibody**

Catalog #: APRab18559



### **Summary**

**Production Name** T2R13 Rabbit Polyclonal Antibody

**Description** Rabbit Polyclonal Antibody

**Host** Rabbit

**Application** IF,WB,ELISA

**Reactivity** Human,Rat,Mouse

#### **Performance**

ConjugationUnconjugatedModificationUnmodified

**Isotype** IgG

ClonalityPolyclonalFormLiquid

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw

cycles.

**Buffer** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.

**Purification** Affinity purification

#### **Immunogen**

Storage

Gene Name TAS2R13

TAS2R13; Taste receptor type 2 member 13; T2R13; Taste receptor family B member 3; **Alternative Names** 

TRB3

**Gene ID** 50838.0

Q9NYV9.The antiserum was produced against synthesized peptide derived from **SwissProt ID** 

human TAS2R13. AA range:123-172

## **Application**

WB 1:500 - 1:2000. IF 1:200 - 1:1000. ELISA: 1:20000. Not yet tested in other

Dilution Ratio

applications.

Molecular Weight 35kD

# **Product Name: T2R13 Rabbit Polyclonal Antibody**

Catalog #: APRab18559




### **Background**

This gene product belongs to the family of candidate taste receptors that are members of the G-protein-coupled receptor superfamily. These proteins are specifically expressed in the taste receptor cells of the tongue and palate epithelia. They are organized in the genome in clusters and are genetically linked to loci that influence bitter perception in mice and humans. In functional expression studies, they respond to bitter tastants. This gene maps to the taste receptor gene cluster on chromosome 12p13. [provided by RefSeq, Jul 2008],function:Receptor that may play a role in the perception of bitterness and is gustducin-linked. May play a role in sensing the chemical composition of the gastrointestinal content. The activity of this receptor may stimulate alpha gustducin, mediate PLC-beta-2 activation and lead to the gating of TRPM5.,miscellaneous:Most taste cells may be activated by a limited number of bitter compounds; individual taste cells can discriminate among bitter stimuli.,similarity:Belongs to the G-protein coupled receptor T2R family.,tissue specificity:Expressed in subsets of taste receptor cells of the tongue and palate epithelium and exclusively in gustducin-positive cells.,

#### **Research Area**

Taste transduction:

## **Image Data**




Immunofluorescence analysis of MCF7 cells, using TAS2R13 Antibody. The picture on the right is blocked with the synthesized peptide.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838


# **Product Name: T2R13 Rabbit Polyclonal Antibody**

Catalog #: APRab18559





Western blot analysis of lysates from Jurkat cells, using TAS2R13 Antibody. The lane on the right is blocked with the synthesized peptide.



Western blot analysis of the lysates from COLO205 cells using TAS2R13 antibody.

#### Note

For research use only.