

Summary

Production Name	VEGF-B Rabbit Polyclonal Antibody
Description	Rabbit Polyclonal Antibody
Host	Rabbit
Application	WB,ELISA
Reactivity	Human,Mouse,Rat

Performance

Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG
Clonality	Polyclonal
Form	Liquid
Storage	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.
Buffer	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% New type preservative N.
Purification	Affinity purification

Immunogen

Gene Name	VEGFB
Alternative Names	VEGFB; VRF; Vascular endothelial growth factor B; VEGF-B; VEGF-related factor; VRF
Gene ID	7423.0
SwissProt ID	P49765.The antiserum was produced against synthesized peptide derived from human VEGFB. AA range:91-140

Application

Dilution Ratio	WB 1:500 - 1:2000. ELISA: 1:20000.
Molecular Weight	21kD

Background

Product Name: VEGF-B Rabbit Polyclonal Antibody
Catalog #: AP Rab19772




This gene encodes a member of the PDGF (platelet-derived growth factor)/VEGF (vascular endothelial growth factor) family. The VEGF family members regulate the formation of blood vessels and are involved in endothelial cell physiology. This member is a ligand for VEGFR-1 (vascular endothelial growth factor receptor 1) and NRP-1 (neuropilin-1). Studies in mice showed that this gene was co-expressed with nuclear-encoded mitochondrial genes and the encoded protein specifically controlled endothelial uptake of fatty acids. Alternatively spliced transcript variants encoding distinct isoforms have been identified. [provided by RefSeq, Sep 2011], alternative products: Additional isoforms seem to exist, function: Growth factor for endothelial cells. VEGF-B167 binds heparin and neuropilin-1 whereas the binding to neuropilin-1 of VEGF-B186 is regulated by proteolysis, PTM: VEGF-B186 is O-glycosylated, similarity: Belongs to the PDGF/VEGF growth factor family, subcellular location: Secreted but remains associated to cells or to the extracellular matrix unless released by heparin, subunit: Homodimer; disulfide-linked. Can also form heterodimer with VEGF, tissue specificity: Expressed in all tissues except liver. Highest levels found in heart, skeletal muscle and pancreas,


Research Area

Cytokine-cytokine receptor interaction; mTOR; Focal adhesion; Pathways in cancer; Renal cell carcinoma; Pancreatic cancer; Bladder cancer;

Image Data




Western blot analysis of lysates from Jurkat cells, using VEGFB Antibody. The lane on the right is blocked with the synthesized peptide.



Western Blot analysis of various cells using VEGF-B Polyclonal Antibody diluted at 1: 2000. Secondary antibody was diluted

Product Name: VEGF-B Rabbit Polyclonal Antibody
Catalog #: APRab19772



Western Blot analysis of Jurkat cells using VEGF-B Polyclonal Antibody diluted at 1 : 2000. Secondary antibody was diluted at 1:20000

Note

For research use only.