
Product Name: GST-Pi Mouse Monoclonal Antibody**Catalog #: AMM21978**

For research use only.

Summary

Description	Mouse Monoclonal Antibody
Host	Mouse
Application	IHC,ELISA
Reactivity	Human,Mouse
Conjugation	Unconjugated
Modification	Unmodified
Isotype	IgG2b,Kappa
Clonality	Monoclonal
Form	Liquid
Storage	Aliquot and store at -20°C (valid for 12 months). Avoid freeze/thaw cycles.
Shipping	Ice bags
Buffer	PBS, 50% glycerol, 0.05% Proclin 300, 0.05%BSA
Purification	The antibody was affinity-purified from ascites by affinity-chromatography using specific immunogen.

Application

Dilution Ratio	IHC 1:200-400;ELISA 1:500-5000
Molecular Weight	Calculated MW:23kDa,Observed MW:23kDa

Antigen Information

Gene Name	GSTP1 FAEES3 GST3 Deafness;Deafness X-linked 7;DFN7;FAEES3;Fatty Acid Ethyl Ester Synthase III;Glutathione S
Alternative Names	Transferase 3;Glutathione S Transferase Pi;Glutathione S-transferase P;Glutathione S-transferase pi 1;GST class-pi;GST3;GSTP;Gstp1;GSTP1-1;GSTP1_HUMAN;PI;X linked 7
Gene ID	Human:2950
SwissProt ID	Human:P09211,Mouse:P19157,Rat:P04906
Immunogen	Synthesized peptide derived from human GST-Pi AA range: 150-210

Background


Glutathione S-transferases (GSTs) are a family of enzymes that play an important role in detoxification by catalyzing the

conjugation of many hydrophobic and electrophilic compounds with reduced glutathione. Based on their biochemical, immunologic, and structural properties, the soluble GSTs are categorized into 4 main classes: alpha, mu, pi, and theta. This GST family member is a polymorphic gene encoding active, functionally different GSTP1 variant proteins that are thought to function in xenobiotic metabolism and play a role in susceptibility to cancer, and other diseases. [provided by RefSeq, Jul 2008],

Research Area

Pathology

Image Data



Human esophageal squamous cell carcinoma tissue was stained with Anti-GST-Pi Antibody