


Product Name: Recombinant Human Artemin
Catalog #: PEH0102



Summary

Name	Artemin
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Artemin is produced by our E.coli expression system and the target gene encoding Ala108-Gly220 is expressed.
Accession #	Q5T4W7
Host	E.coli
Species	Human
Predicted Molecular Mass	12.1 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM HEPES, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human Artemin
Catalog #: PEH0102



Alternative Names

Artemin; ARTN; Enovin; Neublastin; EVN

Background

Human Artemin is a GDNF family ligand that is distantly related to the TGF- β superfamily of molecules. It is synthesized as a preproprotein, and contains a variable length pre-, or signal sequence, plus a 68 amino acid (aa) proregion and a 113 aa mature segment. Following synthesis and proteolytic processing, mature ARTN is secreted as a presumably glycosylated, 28 kDa disulfide-linked homodimer that contains three intrachain disulfide bonds and the typical TGF- β signature cysteine-knot motif. In the mature region, human ARTN is 89% and 88% aa identical to rat and mouse ARTN, respectively. Human ARTN is active on rodent cells. The receptor for ARTN has been identified as the ligand binding subunit GFR α -3 plus the signal transducing subunit, RET. The GFR α -1/RET receptor complex has also been suggested to be a ligand binding unit for ARTN. ARTN is known to be a chemoattractant for sympathetic neuron axons innervating the developing cardiovascular system. It also promotes sensory neuron survival and likely plays a role in the development of the peripheral nervous system. Finally, it has been reported to reverse neuropathic pain due to nerve injury, and to help resolve morphological changes associated with nerve damage.

Note

For Research Use Only , Not for Diagnostic Use.