

Summary

Name	BCL2L1/Bcl-2-like protein 1/B-cell Lymphoma-extra Large/Bcl-xL
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/ μ g as determined by LAL test.
Construction	Recombinant Human B Cell Lymphoma/Leukemia x Long Form is produced by our E.coli expression system and the target gene encoding Met1-Arg212 is expressed with a 6His tag at the C-terminus.
Accession #	Q07817
Host	E.coli
Species	Human
Predicted Molecular Mass	24.85 KDa
Formulation	Supplied as a 0.2 μm filtered solution of 20mM HEPES, 50mM KCl, 20% Glycerol, pH 7.5.
Shipping	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	-

SDS-PAGE image

kDa	MK	R
120 90	-	
60	-	
40	-	
30	-	-
20	-	
14	_	
14		

Background

Alternative Names	Bcl-2-Like Protein 1; Bcl2-L-1; Apoptosis Regulator Bcl-X; BCL2L1; BCL2L; BCLX
Background	Bcl-2-Like Protein 1 (BCL2L1) is a member of the Bcl-2 family. BCL2L1 is expressed at high levels in cells that undergo a high rate of turnover, such as developing



lymphocytes. BCL2L1 is a mitochondrial membrane protein. BCL2L1 contains four motifs, BH1, BH2 and BH4. The BH4 motif is required for anti-apoptotic activity. The BH1 and BH2 motifs are required for both heterodimerization with other Bcl-2 family members and for repression of cell death. BCL2L1 regulates cell death by blocking the voltage-dependent anion channnel (VDAC) and preventing the release of the caspase activator, CYC1, from the mitochondrial membrane. In addition, BCL2L1 promotes apoptosis.

Note

For Research Use Only, Not for Diagnostic Use.