


**Product Name: Recombinant Human CCL26 (71AA)**  
**Catalog #: PEH0266**



## Summary

<b>Name</b>	CCL26/Eotaxin-3 (24-94)
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human C-C Motif Chemokine 26 is produced by our E.coli expression system and the target gene encoding Thr24-Leu94 is expressed.
<b>Accession #</b>	Q9Y258
<b>Host</b>	E.coli
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	8.53 KDa
<b>Formulation</b>	Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 1mM EDTA, 20% Glycerol, pH 9.0.
<b>Shipping</b>	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	0.00.0

## SDS-PAGE image



**Product Name: Recombinant Human CCL26 (71AA)**  
**Catalog #: PEH0266**

---



### **Alternative Names**

C-C Motif Chemokine 26; CC Chemokine IMAC; Eotaxin-3; Macrophage Inflammatory Protein 4-Alpha; MIP-4-Alpha; Small-Inducible Cytokine A26; Thymic Stroma Chemokine-1; TSC-1; CCL26; SCYA26

### **Background**

Chemokine Ligand 26 protein (CCL26) is a novel small cytokine belonging to the CC chemokine family which is involved in immunoregulatory and inflammatory processes. CCL26 is constitutively expressed in thymus, but only transiently expressed in phytohemagglutinin-stimulated peripheral blood mononuclear cells. It specifically binds and induces chemotaxis in T cells and elicits its effects by interacting with the chemokine receptor CCR4. CCL26, along with Eotaxin-1 and Eotaxin-2, selectively activates the CC chemokine receptor 3 (CCR3). The Eotaxin-3-CCR3 interaction may play an important role in allergic diseases such as atopic dermatitis and bronchial asthma. The full-length cDNA for CCL26 encodes a protein of 94 amino acids with a putative signal peptide of either 23 or 26 amino acid residues. Both the 71 and 68 amino acid residue variants of recombinant CCL26 demonstrate equal potency in inducing chemotaxis of a human CCR3-transfected cell line. Unlike most other CC chemokines, CCL26 maps to human chromosome 7q11.2, within 40 kilobases of the Eotaxin-2 loci. CCL26 and Eotaxin-2 are unique in that they are the only chemokines identified to date that map to chromosome 7.

### **Note**

For Research Use Only , Not for Diagnostic Use.