


## Summary

Name	CLIC2/XAP121		
Purity	Greater than 95% as determined by reducing SDS-PAGE		
Endotoxin level	<1 EU/ $\mu$ g as determined by LAL test.		
Construction Accession #	Recombinant Human Chloride Intracellular Channel Protein 2 is produced by our E.coli expression system and the target gene encoding Met1-Ser247 is expressed with a 6His tag at the N-terminus. O15247		
Host	E.coli		
Species	Human		
Predicted Molecular Mass	30.5 KDa		
Predicted Molecular Mass Formulation	30.5 KDa Supplied as a 0.2 μm filtered solution of 20mM Tris-HCl, 100mM NaCl, 1mM DTT, 20% Glycerol, pH 8.0.		
	Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 100mM NaCl, 1mM DTT, 20% Glycerol, pH 8.0. The product is shipped on dry ice/polar packs. Upon receipt, store it immediately		
Formulation	Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 100mM NaCl, 1mM DTT, 20% Glycerol, pH 8.0.		

## **SDS-PAGE** image



## Background

Alternative Names	Chloride Intracellular Channel Protein 2; XAP121; CLIC2			
Background	Chloride Intracellular Channel Protein 2 (CLIC2) is a critical component of all living			
Web: https://www.enkilife.com	E-mail: order@enkilife.com	techsupport@enkilife.com	Tel: 0086-27-87002838	



cells; it regulatescellular traffic of Chloride ion and it can be inserted into membranes and form chloride ion channels. Membrane insertion seems to be redox-regulated and may occur only under oxydizing conditions, channel activity depends on the pH. CLIC2 is involved in regulating membrane potential and organic solute transport. CLIC2 modulates the activity of RYR2 and inhibits Calcium influx. CLIC2 can be detected in the adult brain, liver, lung, heart, stomach, spleen and testis. It is expressed in fetal liver and adult skeletal muscle. CLIC2 is a potential candidate for one of many diseases linked to Xq28.

## Note

For Research Use Only, Not for Diagnostic Use.