

Product Name: Recombinant Human Cornulin (N-6His)
Catalog #: PEH0446




Summary

Name	Cornulin
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Cornulin is produced by our E.coli expression system and the target gene encoding Met1-Ser140 is expressed with a 6His tag at the N-terminus.
Accession #	Q9UBG3
Host	E.coli
Species	Human
Predicted Molecular Mass	17.45 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human Cornulin (N-6His)
Catalog #: PEH0446



Alternative Names

Cornulin; 53 kDa Putative Calcium-Binding Protein; 53 kDa Squamous Epithelial-Induced Stress Protein; 58 kDa Heat Shock Protein; Squamous Epithelial Heat Shock Protein 53; Tumor-Related Protein; CRNN; C1orf10; DRC1; PDRC1; SEP53

Background

Cornulin is a member of the fused gene family of molecular chaperones. Human Cornulin contains N-terminus EF-hand domains and Ca^{2+} binding domains, and two glutamine- and threonine-rich 60 amino acid repeats in its C-terminus. Cornulin involves in the mucosal/epithelial immune response and epidermal differentiation. Cornulin is a survival factor that participates in the clonogenicity of squamous esophageal epithelium cell lines, attenuates deoxycholic acid (DCA)-induced apoptotic cell death and release of calcium. When Cornulin is overexpressed in oral squamous carcinoma cell lines, it regulates negatively cell proliferation by the induction of G1 arrest.

Note

For Research Use Only , Not for Diagnostic Use.