


Product Name: Recombinant Human IL-8 (72AA)
Catalog #: PEH0482



Summary

Name	CXCL8/IL-8 (8-79)(72aa)(Ser28-Ser99)
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Interleukin-8 is produced by our E.coli expression system and the target gene encoding Ser28-Ser99 is expressed.
Accession #	P10145
Host	E.coli
Species	Human
Predicted Molecular Mass	8.45 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human IL-8 (72AA)
Catalog #: PEH0482

Alternative Names

Interleukin-8; IL-8; C-X-C Motif Chemokine 8; Emotakin; Granulocyte Chemotactic Protein 1; GCP-1; Monocyte-Derived Neutrophil Chemotactic Factor; MDNCF; Monocyte-Derived Neutrophil-Activating Peptide; MONAP; Neutrophil-Activating Protein 1; NAP-1; Protein 3-10C; T-Cell Chemotactic Factor; IL8; CXCL8

Background

Interleukin-8 (IL-8) belongs to the neutrophil-specific CXC family of chemokines. It is one of the initial cytokines released from a variety of cell types, including T cells, endothelial cells and fibroblasts, in response to an inflammatory stimulus and acts by recruiting neutrophils, T-cells and basophils to the site of inflammation. Elevated Interleukin-8 levels are associated with the onset of a variety of disease states.

Note

For Research Use Only , Not for Diagnostic Use.