Product Name: Recombinant Human CSTA (N-6His)

Catalog #: PEH0491



Summary

Name Cystatin A

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Cystatin A is produced by our E.coli expression system

and the target gene encoding Ile2-Phe98 is expressed with a 6His tag at the

N-terminus.

Accession # P01040

Host E.coli

Species Human

Predicted Molecular Mass 11.8 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 100mM NaCl, pH

8.0.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability & Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human CSTA (N-6His)

Catalog #: PEH0491



Alternative Names Cystatin-A; Cystatin-AS; Stefin-A; CSTA; STF1; STFA

Background Human Cystatin A (CSTA) is a member of family 1 of the cystatin superfamily,

which is characterized by lacking of disulphide bonds an carbohydrates. Cystatin A is an intracellular inhibitor regulating the activities of cysteine proteases of the papain family such as Cathepsins B, H and L. Cystatin A is also implicated in a number of disease states. Due to altered proteolytic state in cancer progression,

Cystatin A may play a role in the proteolytic pathways.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838