


**Product Name: Recombinant Human EIF1AX (N-6His)**  
**Catalog #: PEH0562**



## Summary

<b>Name</b>	EIF1AX
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human Eukaryotic Translation Initiation Factor 1A, X-Chromosomal is produced by our E.coli expression system and the target gene encoding Met1-Ile144 is expressed with a 6His tag at the N-terminus.
<b>Accession #</b>	P47813
<b>Host</b>	E.coli
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	18.6 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image



## Background

**Product Name: Recombinant Human EIF1AX (N-6His)**  
**Catalog #: PEH0562**

---



**Alternative Names**

Eukaryotic Translation Initiation Factor 1A X-Chromosomal; eIF-1A X Isoform; Eukaryotic Translation Initiation Factor 4C; eIF-4C; EIF1AX; EIF1A; EIF4C

**Background**

Eukaryotic Translation Initiation Factor 1A, X-Chromosomal (EIF1AX) is an essential eukaryotic translation initiation factor that belongs to the eIF-1A family. EIF1AX is required for the binding of the 43S complex (a 40S subunit, eIF2/GTP/Met-tRNA<sub>i</sub> and eIF3) to the 5' end of capped RNA and has been shown to interact with IPO13. EIF1AX contains one S1-like domain and seems to be required for maximal rate of protein biosynthesis. Enhances ribosome dissociation into subunits and stabilizes the binding of the initiator Met-tRNA(I) to 40 S ribosomal subunits.

**Note**

For Research Use Only , Not for Diagnostic Use.