


## Summary

Name	FABP5/E-FABP
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Human Fatty Acid-Binding Protein 5 is produced by our E.coli expression system and the target gene encoding Ala2-Glu135 is expressed with a 6His tag at the N-terminus.
Accession #	Q01469
Host	E.coli
Species	Human
Predicted Molecular Mass	17.33 KDa
Formulation	Lyophilized from a 0.2 $\mu$ m filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at $\leq$ -70°C, stable for 6 months after receipt. Store at $\leq$ -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## **SDS-PAGE** image



## Background



Alternative NamesFatty Acid-Binding Protein Epidermal; Epidermal-Type Fatty Acid-Binding Protein;<br/>E-FABP; Fatty Acid-Binding Protein 5; Psoriasis-Associated Fatty Acid-Binding<br/>Protein Homolog; PA-FABP; FABP5BackgroundFatty acid-binding protein 5 (FABP5) is a cytoplasm protein that belongs to the<br/>fatty-acid binding protein (FABP) family of calycin superfamily. Fatty acid binding<br/>proteins are a family of small, highly conserved, cytoplasmic proteins that bind<br/>long-chain fatty acids. FABP5 can be expressed in keratinocytes, and is highly<br/>expressed in psoriatic skin. FABP5 has been shown to be involved in keratinocyte<br/>differentiation. FABP5 has high specificity for fatty acids, the highest affinity for<br/>C18 chain length. FABP5 can decrease the chain length or introduce double bonds<br/>to reduce the affinity.

## Note

For Research Use Only, Not for Diagnostic Use.