## **Product Name: Recombinant Human Grancalcin**

Catalog #: PEH0756



#### **Summary**

Name Grancalcin/GCA

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/μg as determined by LAL test.

Construction Recombinant Human Grancalcin is produced by our E.coli expression system

and the target gene encoding Met1-Ile217 is expressed.

Accession # P28676

**Host** E.coli

**Species** Human

Predicted Molecular Mass 24.01 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, 1mM

EDTA, pH 8.5.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

**Stability&Storage** Store at  $\leq$ -70°C, stable for 6 months after receipt. Store at  $\leq$ -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

**Reconstitution** Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

#### **SDS-PAGE** image



### **Background**

# **Product Name: Recombinant Human Grancalcin Catalog #: PEH0756**



Alternative Names Grancalcin; GCA; GCL

Background Grancalcin (GCA) is a cytoplasmic granule membrane protein that contains 4 EF-

hand domains. GCA is calcium-binding protein and particularly abundant in human neutrophils. GCA is highly expressed in bone marrow, and it can be detected in neutrophils and macrophages. Calcium-binding protein GCA cooperates with SRI and LCP1, so it may play a role in the adhesion of neutrophils to fibronectin. GCA

also may play a role in the formation of focal adhesions.

#### Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838