EnkiLife Product Name: Recombinant Human ASF1A (C-6His, N-T7 tag) Catalog #: PEH0794

Summary

Name Histone chaperone ASF1A

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/µg as determined by LAL test.

Construction Recombinant Human Histone Chaperone ASF1A is produced by our E.coli

expression system and the target gene encoding Met1-Met204 is expressed

with a T7 tag at the N-terminus, 6His tag at the C-terminus.

Accession # O9Y294

Host E.coli **Species**

Predicted Molecular Mass 25.4 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 1mM DTT, 150mM

NaCl, pH 8.0.

Human

The product is shipped at ambient temperature. Upon receipt, store it **Shipping**

immediately at the temperature listed below.

Stability&Storage Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw


cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838



Alternative Names

Histone Chaperone ASF1A; Anti-Silencing Function Protein 1 Homolog A; hAsf1; hAsf1a; CCG1-Interacting Factor A; CIA; hCIA; ASF1A

Background

Human Histone Chaperone ASF1A (ASF1A) belongs to the H3/H4 family of histone chaperone proteins. ASF1A is ubiquitously expressed in many cells and tissues, interacting with histones H3 and H4. ASF1A cooperates with Chromatin Assembly Factor 1 to promote replication-dependent chromatin assembly and with HIRA to promote replication-independent chromatin assembly. In addition, ASF1A is necessary for the formation of senescence-associated heterochromatin foci (SAHF) and efficient senescenceassociated cell cycle exit.

Note

For Research Use Only, Not for Diagnostic Use.