

Summary

Name HMGB1/High mobility group protein B1/HMG-1(Truncated)

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human High Mobility Group Protein B1 is produced by our

E.coli expression system and the target gene encoding Pro92-Val176 is

expressed.

Accession # P09429

Host E.coli

Species Human

Predicted Molecular Mass 10 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 50mM HEPES-Na, 500mM NaCl,

0.6mM DTT, pH 7.9.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt.


Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at \leq -20°C for 3 months.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human HMGB1 (Truncated) Enkilife Catalog #: PEH0799

Alternative Names High Mobility Group Protein B1; High Mobility Group Protein 1; HMG-1; HMGB1;

HMG1

Background High mobility group protein B1 is a member of the HMGB family consisting of

three members, HMGB1, HMGB2 and HMGB3.It Contains 2 HMG box DNA-binding domains entitled box A and box B and It is a highly negative-charged C terminus. As a nuclear protein, HMGB1 stabilizes nucleosomes and allows bending of DNA that facilitates gene transcription which is essential for individual survival. Meanwhile, it is revealed that HMGB1 can also act as a cytokine extracellularly and regulates monocyte, T cell, dendritic cell activities in inflammatory responses.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838