Product Name: Recombinant Human IGF-I (67AA)
Catalog #: PEH0839

Summary

Name	IGF-I
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<0.5 EU/μg as determined by LAL test.
Construction	Recombinant Human Insulin-like Growth Factor I(4-70) is produced by our E.coli expression system and the target gene encoding Thr52-Ala118 is expressed.
Accession #	P05019
Host	E.coli
Species	Human
Predicted Molecular Mass	7.3 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM NaAc-HAc, pH 4.5
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human IGF-I (67AA)
Catalog #: PEH0839

Alternative Names	Insulin-Like Growth Factor I; IGF-I; Mechano Growth Factor; MGF; Somatomedin-C; IGF1; IBP1
Background	Insulin-like growth factor I (IGF1) belongs to the family of insulin-like growth factors that are structurally homologous to proinsulin. Mature IGFs are generated by proteolytic processing of inactive precursor protein containing N-terminal and C-terminal propeptide regions. Mature human IGF-I consisting of 70 amino acids with 94% identity with mouse IGF1 and exhibits cross-species activity. IGF1 binds IGF-1R, IGF-2R, and the insulin receptor and plays a key role in cell cycle progression, cell proliferation and tumor progression. IGF1 expression is regulated by growth hormone.

Note

For Research Use Only , Not for Diagnostic Use.