


Product Name: Recombinant Human IL-1a
Catalog #: PEH0888



Summary

Name	IL-1 alpha/IL-1F1/IL-1 α /IL-1a/Interleukin-1 Alpha
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<0.01 EU/ μ g as determined by LAL test.
Construction	Recombinant Human Interleukin-1 Alpha is produced by our E.coli expression system and the target gene encoding Ser113-Ala271 is expressed.
Accession #	P01583
Host	E.coli
Species	Human
Predicted Molecular Mass	18 KDa
Formulation	Lyophilized from a 0.2 μ m filtered solution of 20mM Tris-HCl, 150mM NaCl, pH 7.5.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at $\leq -20^{\circ}\text{C}$, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8 $^{\circ}\text{C}$ for 2-7 days. Aliquots of reconstituted samples are stable at $\leq -20^{\circ}\text{C}$ for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100 μ g/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human IL-1a
Catalog #: PEH0888

Alternative Names

Interleukin-1 Alpha; IL-1 Alpha; Hematopoietin-1; IL1A; IL1F1

Background

Interleukin-1 alpha (IL1 α) is a cytokine member of the interleukin-1 family. IL-1 consists of two distinct forms: IL1 α and IL1 β that recognize the same cell surface receptors but are distinct proteins with approximately 25% amino acid sequence identity. IL1 α is constitutively produced by epithelial cells and plays an essential role in maintenance of skin barrier function. Upon stimulation, a wide variety of cells including osteoblasts, monocytes, macrophages can be induced to express IL1 α . IL1 α possesses a wide range of metabolic, physiological, haematopoietic activities, and is critically involved in the regulation of the immune responses and inflammatory responses.

Note

For Research Use Only , Not for Diagnostic Use.