


Summary

Name	ISG15/G1P2/UCRP
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Interferon-stimulated Gene 15 is produced by our E.coli expression system and the target gene encoding Gly2-Gly157 is expressed with a 6His tag at the C-terminus.
Accession #	AAH09507.1
Host	E.coli
Species	Human
Predicted Molecular Mass	18.1 KDa
Formulation	Supplied as a 0.2 μm filtered solution of 50mM HEPES, 100mM NaCl, pH 8.0.
Shipping	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	

SDS-PAGE image



Alternative Names

Ubiquitin-Like Protein ISG15; Interferon-Induced 15 kDa Protein; Interferon-Induced 17 kDa Protein; IP17; Ubiquitin Cross-Reactive Protein; hUCRP; ISG15; G1P2; UCRP

Background

Ubiquitin-Like Protein ISG15 (ISG15) is a ubiquitin-like protein that becomes conjugated to many cellular proteins upon activation by interferon-alpha and -beta. Several functions have been ascribed to the encoded protein, including chemotactic activity towards neutrophils, direction of ligated target proteins to intermediate filaments, cell-to-cell signaling, and antiviral activity during viral infections. While conjugates of this protein have been found to be noncovalently attached to intermediate filaments, this protein is sometimes secreted. ISG15 becomes conjugated to a diverse set of proteins after IFN-alpha/beta stimulation or microbial challenge. The functions or biochemical consequences ISG15 conjugation to proteins are not yet known, but it appears that this modification does not target proteins for proteasomal degradation. ISG15 shows specific chemotactic activity towards neutrophils and activates them to induce release of eosinophil chemotactic factors. Upon interferon treatment, ISG15 can be detected in both free and conjugated forms, and is secreted from monocytes and lymphocytes where it can function as a cytokine. In the cell, ISG15 co-localizes with intermediate filaments and ISGylation may modulate the JAK-STAT pathway or certain aspects of neurological disease.

Note

For Research Use Only, Not for Diagnostic Use.