Product Name: Recombinant Human MBIPP (N-6His) Catalog #: PEH1137



Summary

Name MBIP/MAP3K12-binding inhibitory protein 1

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human MAP3K12-Binding Inhibitory Protein 1 is produced by

our E.coli expression system and the target gene encoding Met1-Pro344 is

expressed with a 6His tag at the N-terminus.

Accession # Q9NS73

Host E.coli

Species Human

Predicted Molecular Mass 42.51 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, pH 8.0.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human MBIPP (N-6His) Catalog #: PEH1137



Alternative Names

MAP3K12-Binding Inhibitory Protein 1; MAPK Upstream Kinase-Binding Inhibitory

Protein; MUK-Binding Inhibitory Protein; MBIP

Background

MAP3K12-binding inhibitory protein 1 (MBIP) is a 39kD protein high expression in the heart and lung. It is a component of the ADA2A-containing complex (ATAC) complex, a complex with histone acetyltransferase activity on histones H3 and H4, and composed of CSRP2BP, KAT2A, TADA2L, TADA3L, ZZ3, MBIP, WDR5, YEATS2, CCDC101 and DR1. In the complex, it probably interacts directly with KAT2A, CSRP2BP and WDR5. It's function to inhibit the MAP3K12 activity to induce the

activation of the JNK/SAPK pathway.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838