


Summary

Name	Nucleobindin-2/Nesfatin-1
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Human Nucleobindin-2 is produced by our E.coli expression
Accession #	system and the target gene encoding Val25-Leu106 is expressed. P80303
Host	E.coli
Species	Human
Predicted Molecular Mass	9.6 KDa
Formulation	Luch his address and a sum filtered colution of 10mM Sodium Dhosenhote all C.E.
Formulation	Lyophilized from a 0.2 μ m filtered solution of 10mM Sodium Phosphate, pH 6.5.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it

SDS-PAGE image



Background



Alternative NamesNucleobindin-2; DNA-binding protein NEFA; Gastric cancer antigen Zg4;
Prepronesfatin; Nesfatin-1; NUCB2; NEFABackgroundNesfatin-1 is a metabolic polypeptide encoded in the N-terminal region of the
precursor protein, Nucleobindin2 (NUCB2). Nesfatin-1 is a neuropeptide produced
in the hypothalamus of mammals. It participates in the regulation of hunger and
fat storage. Nesfatin-1 is also expressed in other areas of the brain, and in
pancreatic islets β -cells, gastric endocrine cells and adipocytes. Nesfatin-1
suppresses food intake and can regulate energy metabolism in a Leptin
independent manner. Nesfatin-1 may also exert hypertensive roles and modulate
blood pressure through directly acting on peripheral arterial resistance.

Note

For Research Use Only, Not for Diagnostic Use.