Catalog #: PEH1257



Summary

CDKN1B/Cyclin-dependent kinase inhibitor 1B/p27/Kip1 Name

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/µg as determined by LAL test.

Construction Recombinant Human Cyclin-Dependent Kinase Inhibitor 1B is produced by

our E.coli expression system and the target gene encoding Met1-Thr198 is

expressed with a 6His tag at the N-terminus.

Accession # P46527

Host E.coli

Species Human

Predicted Molecular Mass 24.2 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 Stability&Storage

months under sterile conditions after opening. Please minimize freeze-thaw

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is


not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838



Catalog #: PEH1257



Alternative Names

Cyclin-Dependent Kinase Inhibitor 1B; Cyclin-Dependent Kinase Inhibitor p27; p27Kip1; CDKN1B; KIP1

Background

Cyclin-Dependent Kinase Inhibitor 1B (CDKN1B) is a Kinesin-related motor protein necessary for mitotic spindle assembly and chromosome segregation. CDKN1B is expressed in all tissues with highest levels observed in skeletal muscle. CDKN1B is a potent inhibitor of Cyclin E- and Cyclin A-CDK2 complexes. CDKN1B forms a complex with Cyclin Type D-CDK4 complexes and is involved in the assembly, stability, and modulation of CCND1-CDK4 complex activation. In addition, CDKN1B acts as an inhibitor or an activator of Cyclin Type D-CDK4 complexes depending on its phosphorylation state and stoichometry.

Note

For Research Use Only, Not for Diagnostic Use.