Product Name: Recombinant Human PDCD10

Catalog #: PEH1279



Summary

Name PDCD10/Programmed cell death protein 10

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Programmed Cell Death Protein 10 is produced by our

E.coli expression system and the target gene encoding Met1-Ala212 is

expressed.

Accession # Q9BUL8

Host E.coli

Species Human

Predicted Molecular Mass 24.9 KDa

Formulation Lyophilized from a 0.2 μm filtered solution of 25mM Tris-HCl, pH 7.3.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.


SDS-PAGE image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

Product Name: Recombinant Human PDCD10

Catalog #: PEH1279





Alternative Names

Programmed Cell Death Protein 10; Cerebral Cavernous Malformations 3 Protein; TF-1 Cell Apoptosis-Related Protein 15; PDCD10; CCM3; TFAR15

Background

Programmed Cell Death Protein 10 (PDCD10) belongs to the PDCD10 family. PDCD10 exists as a homodimer and is widely expressed. PDCD10 can increase mitogen-activated protein kinase activity and MST4 activity. PDCD10 is required for normal cardiovascular development and normal angiogenesis, vasculogenesis and hematopoiesis during embryonic development. Defects in PDCD10 are the cause of cerebral cavernous malformations type 3.

Note

For Research Use Only, Not for Diagnostic Use.