


**Product Name: Recombinant Human PDCD4 (C-6His)**  
**Catalog #: PEH1280**



## Summary

<b>Name</b>	PDCD4/Programmed cell death protein 4
<b>Purity</b>	Greater than 95% as determined by reducing SDS-PAGE
<b>Endotoxin level</b>	<1 EU/μg as determined by LAL test.
<b>Construction</b>	Recombinant Human Programmed Cell Death Protein 4 is produced by our E.coli expression system and the target gene encoding Lys212-Pro357 is expressed with a 6His tag at the C-terminus.
<b>Accession #</b>	Q53EL6
<b>Host</b>	E.coli
<b>Species</b>	Human
<b>Predicted Molecular Mass</b>	17 KDa
<b>Formulation</b>	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
<b>Shipping</b>	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
<b>Stability&amp;Storage</b>	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
<b>Reconstitution</b>	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

## SDS-PAGE image



**Product Name: Recombinant Human PDCD4 (C-6His)**  
**Catalog #: PEH1280**

---



## Background

### Alternative Names

Programmed Cell Death Protein 4; Neoplastic Transformation Inhibitor Protein; Nuclear Antigen H731-Like; Protein 197/15a; PDCD4; H731

### Background

Programmed Cell Death Protein 4 (PDCD4) is a member of the PDCD4 family. PDCD4 and EIF4A1 form a heterotrimer. One molecule of PDCD4 binds two molecules of EIF4A1. PDCD4 takes part in apoptosis via inhibiting translation initiation and cap-dependent translation. PDCD4 promotes colonic neoplastic transformation and tumor invasion. PDCD4 is an important target for microRNA R-21 in breast cancer cells. Shortage of PDCD4 expression is associated with colorectal cancer. Overexpression of PDCD4 in carcinoid cells results in inhibition of cell proliferation.

## Note

For Research Use Only , Not for Diagnostic Use.