


Summary

Name	PDGF-BB
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<0.2 EU/μg as determined by LAL test.
Construction	Recombinant Human Platelet-Derived Growth Factor BB is produced by our E.coli expression system and the target gene encoding Ser82-Thr190 is expressed.
Accession #	P01127
Host	E.coli
Species	Human
Predicted Molecular Mass	12.42 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM NaAc-HAc, pH 4.5.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human PDGF-BB
Catalog #: PEH1285



Alternative Names

PDGFBB; PDGF-BB

Background

Platelet-Derived Growth Factor Subunit B (PDGFB) belongs to the PDGF/VEGF growth factor family. Platelet-derived growth factor is a potent mitogen for cells of mesenchymal origin. PDGFB can exist either as a homodimer (PDGF-BB) or as a heterodimer with the platelet-derived growth factor alpha polypeptide (PDGF-AB), where the dimers are connected by disulfide bonds. Mutations in this gene are associated with meningioma. Binding of PDGFB to its receptor elicits a variety of cellular responses. In addition, PDGFB is released by platelets upon wounding and plays an important role in stimulating adjacent cells to grow and thereby heals the wound.

Note

For Research Use Only , Not for Diagnostic Use.