

Product Name: Recombinant Human PEA15
Catalog #: PEH1298




Summary

Name	PEA15/PED/Astrocytic phosphoprotein PEA-15
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Phosphoprotein Enriched in Astrocytes of 15 kDa is produced by our E.coli expression system and the target gene encoding Met1-Ala130 is expressed.
Accession #	Q15121
Host	E.coli
Species	Human
Predicted Molecular Mass	15.3 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human PEA15
Catalog #: PEH1298



Alternative Names

Astrocytic Phosphoprotein PEA-15; 15 kDa Phosphoprotein Enriched in Astrocytes; Phosphoprotein Enriched in Diabetes; PED; PEA15

Background

Astrocytic phosphoprotein PEA-15 (PEA15) is a death effector domain (DED)-containing protein. PEA15 is mainly expressed in the central nervous system, principally in astrocytes. Increased PEA15 levels affect tumorigenesis and cancer progression. PEA15 is overexpressed in breast cancers and gliomas as well as in type 2 diabetes. PEA15 blocks Ras-mediated inhibition of integrin activation and modulates the ERK MAP kinase cascade. PEA15 also inhibits RPS6KA3 activities by holding it in the cytoplasm. In addition, PEA15 inhibits both TNFRSF6 and TNFRSF1A mediated CASP8 activity and apoptosis. At present, PEA15 expression is also a significant prognostic marker in ovarian cancer.

Note

For Research Use Only , Not for Diagnostic Use.