


Summary

Name	Peptidyl-Prolyl Cis-Trans Isomerase H/Ppih
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/µg as determined by LAL test.
Construction	Recombinant Human Peptidyl-Prolyl Cis-Trans Isomerase H is produced by our E.coli expression system and the target gene encoding Met1-Met177 is expressed with a 6His tag at the N-terminus.
Accession #	O43447
Host	E.coli
Species	Human
Predicted Molecular Mass	21.4 KDa
Formulation	Supplied as a 0.2 μ m filtered solution of PBS, pH 7.4.
Shipping	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	

SDS-PAGE image



Background

Alternative Names Pep	tidyl-Prolyl Cis-Trans Isomerase H; PPlase H; Rotamase H; Small Nuclear
Rib	onucleoprotein Particle-Specific Cyclophilin H; CypH; U-snRNP-Associated
Сус	lophilin SnuCyp-20; USA-CYP; PPIH; CYP20; CYPH
Background Pep	tidyl-Prolyl Cis-Trans Isomerase H (PPIH) belongs to the Cyclophilin-type
PPI	ase family that accelerate the folding of proteins. PPlases can catalyze the cis-

Product Name: Recombinant Human PPIH (N-6His) Catalog #: PEH1305



trans isomerization of Proline Imidic peptide bonds in oligopeptides. PPIH participates in pre-mRNA splicing. It is a specific component of the complex that includes pre-mRNA processing factors PRPF3, PRPF4, and PRPF18, as well as U4/U5/U6 tri-snRNP. In addition, PPIH has PPIase activity and may play a role as a chaperone mediating the interactions between different proteins inside the spliceosome.

Note

For Research Use Only, Not for Diagnostic Use.