

Summary

Name	Prefoldin subunit 2/PFDN2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/ μ g as determined by LAL test.
Construction	Recombinant Human Prefoldin Subunit 2 is produced by our E.coli expression system and the target gene encoding Met1-Ser154 is expressed with a 6His tag at the N-terminus.
Accession #	Q9UHV9
Host	E.coli
Species	Human
Predicted Molecular Mass	18.8 KDa
Formulation	Lyophilized from a 0.2 μ m filtered solution of 20mM Tris-HCl, 50mM NaCl, 1mM DTT, pH 8.0.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image





Background

Alternative NamesPrefoldin Subunit 2; PFDN2; PFD2BackgroundPrefoldin Subunit 2 (PFDN2) belongs to the Prefoldin Beta subunit family. The
PFDN2 protein is one of six subunits of Prefoldin that act as a molecular chaperone
complex that binds and stabilizes newly synthesized polypeptides allowing them to
fold correctly. PFDN2 binds specifically to Cytosolic Chaperonin (c-CPN) and
transfers target proteins to it. PFDN2 also binds to a nascent polypeptide chain
and promotes folding in settings where there are many competing pathways for
non-native proteins.

Note

For Research Use Only, Not for Diagnostic Use.