

Product Name: Recombinant Human S100A13
Catalog #: PEH1390




Summary

Name	Protein S100-A13/S100A13
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Protein S100-A13 is produced by our E.coli expression system and the target gene encoding Ala2-Lys98 is expressed.
Accession #	Q99584
Host	E.coli
Species	Human
Predicted Molecular Mass	11.3 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 50mM Tris-HCl, 1mM CaCl ₂ , 0.1% Tween-20, pH8.0.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human S100A13
Catalog #: PEH1390



Alternative Names

Protein S100-A13;S100A13;S100 calcium-binding protein A13;

Background

S100A13 is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs. It is widely expressed in various types of tissues with a high expression level in thyroid gland. In smooth muscle cells, this protein co-expresses with other family members in the nucleus and in stress fibers, suggesting diverse functions in signal transduction. It plays a role in the export of proteins that lack a signal peptide and are secreted by an alternative pathway. It binds two calcium ions per subunit and one copper ion. Binding of one copper ion does not interfere with calcium binding. It is required for the copper-dependent stress-induced export of IL1A and FGF1. The calcium-free protein binds to lipid vesicles containing phosphatidylserine, but not to vesicles containing phosphatidylcholine.

Note

For Research Use Only , Not for Diagnostic Use.