


Product Name: Recombinant Human RBP5
Catalog #: PEH1434



Summary

Name	RBP5/Retinol-binding Protein 5
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Retinol-binding Protein 5 is produced by our E.coli expression system and the target gene encoding Met1-Arg135 is expressed.
Accession #	P82980
Host	E.coli
Species	Human
Predicted Molecular Mass	15.9 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human RBP5
Catalog #: PEH1434

Alternative Names

Retinol-binding protein 5; Cellular retinol-binding protein III; CRBP-III; HRBPiso; RBP5

Background

Retinol-binding proteins (RBP) are a family of proteins with diverse functions. They are carrier proteins that bind retinol. Retinol and retinoic acid play crucial roles in the modulation of gene expression and overall development of an embryo. However, deficit or excess of either one of these substances can cause early embryo mortality or developmental malformations. Regulation of transport and metabolism of retinol necessary for a successful pregnancy is accomplished via RBP. Retinol binding proteins have been identified within the uterus, embryo, and extraembryonic tissue of the bovine, ovine, and porcine, clearly indicating that RBP plays a role in proper retinol exposure to the embryo and successful transport at the maternal-fetal interface.

Note

For Research Use Only , Not for Diagnostic Use.