

Product Name: Recombinant Human RheB (N-GST)
Catalog #: PEH1438




Summary

Name	RHEB/RHEB2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Human Ras Homolog Enriched In Brain is produced by our E.coli expression system and the target gene encoding Met1-Met184 is expressed with a GST tag at the N-terminus.
Accession #	Q15382
Host	E.coli
Species	Human
Predicted Molecular Mass	20.4 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of 20mM PB, 8% Trehalose, 4% Mannitol, 50mM NaCl, 10mM GSH, 0.05% Tween 80, pH6.5.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Human RheB (N-GST)
Catalog #: PEH1438



Alternative Names

GTP-Binding Protein Rheb; Ras Homolog Enriched in Brain; RHEB; RHEB2

Background

GTP-Binding Protein Rheb (RHEB) is a member of the small GTPase superfamily and encodes a lipid-anchored, cell membrane protein with five repeats of the RAS-related GTP-binding region. Highest levels of RHEB can be found in the skeletal and cardiac muscle, and it is vital in the regulation of growth and cell cycle progression due to its role in the Insulin/TOR/S6K signaling pathway. RHEB stimulates the phosphorylation of S6K1 and EIF4EBP1 through activation of mTORC1 signaling, and it activates the protein kinase activity of mTORC1. RHEB has GTPase activity and shuttles between a GDP-bound form and a GTP-bound form, farnesylation of the protein is required for this activity.

Note

For Research Use Only , Not for Diagnostic Use.