# Catalog #: PEH1523



### **Summary**

Name SH2D1A/SH2 domain-containing protein 1A

**Purity** Greater than 95% as determined by reducing SDS-PAGE

**Endotoxin level** <1 EU/µg as determined by LAL test.

Human

Construction Recombinant Human SH2 Domain-Containing Protein 1A is produced by our

E.coli expression system and the target gene encoding Met1-Pro128 is

expressed with a 6His tag at the N-terminus.

Accession # O60880

Host E.coli **Species** 

**Predicted Molecular Mass** 16.4 KDa

**Formulation** Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, 10%

Glycerol, pH 7.5.

The product is shipped on dry ice/polar packs. Upon receipt, store it immediately **Shipping** 

at the temperature listed below.

Stability&Storage Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3

months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution

**SDS-PAGE** image

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838

# **Product Name: Recombinant Human SH2D1A (N-6His)** Catalog #: PEH1523

**C** EnkiLife

#### **Alternative Names**

SH2 Domain-Containing Protein 1A; Duncan Disease SH2-Protein; Signaling Lymphocytic Activation Molecule-Associated Protein; SLAM-Associated Protein; T-Cell Signal Transduction Molecule SAP; SH2D1A; DSHP; SAP

## **Background**

SH2 Domain-Containing Protein 1A,(SH2D1A) contains one SH2 domain and a short tail and is localized in the cytoplasm. SH2D1A is expressed at a high level in the thymus and the lung, with lower expression levels in the spleen and the liver. SH2D1A acts as an inhibitor of the signaling lymphocyte activation molecule (SLAM) self-association. In addition, SH2D1A mediates interaction between FYN and SLAMF1. It is also thought to regulate the activity of the neurotrophin receptors NTRK1, NTRK2 and NTRK3.

#### Note

For Research Use Only, Not for Diagnostic Use.