Product Name: Recombinant Human STAT3 (C-6His)

Catalog #: PEH1568



Summary

Name STAT3

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Signal Transducer And Activator of Transcription 3 is

produced by our E.coli expression system and the target gene encoding

Met1-Asn175 is expressed with a 6His tag at the C-terminus.

Accession # P40763

Host E.coli

Species Human

Predicted Molecular Mass 21.8 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, 10% Trehalose,

50mM NaCl, 0.05% Tween 80, pH 8.0.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human STAT3 (C-6His) Catalog #: PEH1568



Alternative Names Signal Transducer and Activator of Transcription 3; Acute-Phase Response Factor;

STAT3; APRF

Background Signal Transducer and Activator of Transcription 3 (STAT3) belongs to the

transcription factor STAT family. STAT3 contains one SH2 domain and is a transcription factor expressed in most cell types. STAT3 is activated by multiple cytokines and growth factors including: IFN-a, IL-10, IL-6, IL-11, IL-12, IL-2, EGF etc. STAT3 functions as signal transducer and transcription activator that mediates cellular responses to interleukins, KITLG/SCF and other growth factors. In addition, STAT3 may also mediate cellular responses to activated FGFR1, FGFR2, FGFR3 and

FGFR4.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838