Product Name: Recombinant Human TAC3 (N-6His)

Catalog #: PEH1603



Summary

Name Tachykinin-3/TAC3

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Human Tachykinin-3 is produced by our E.coli expression

system and the target gene encoding Gln17-Glu121 is expressed with a 6His

tag at the N-terminus.

Accession # Q9UHF0

Host E.coli

Species Human

Predicted Molecular Mass 13.9 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM Tris-HCl, pH 8.0.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3


months under sterile conditions after opening. Please minimize freeze-thaw

cycles.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Human TAC3 (N-6His)

Catalog #: PEH1603



Alternative Names

Tachykinin-3; ZNEUROK1; Neurokinin-B; NKB; Neuromedin-K; TAC3; NKNB; UNQ585/PRO1155

Background

Tachykinin 3 (TAC3) is a secreted protein that belongs to the Tachykinin family. Tachykinins are active peptides that excite neurons and evoke behavioral responses; they are potent vasodilators and secretagogues, and contract many smooth muscles in pregnancy. TAC3 is primarily expressed in the central and peripheral nervous systems and functions as a neurotransmitter. It is also expressed in the outer syncytiotrophoblast of the placenta and may be associated with pregnancy-induced hypertension and pre-eclampsia. TAC3 acts as the ligand for the neurokinin-3 receptor, mutations in this gene are associated with normosmic hypogonadotropic hypogonadism.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838