


Summary

| Name | Adiponectin/Acrp30/ADIPOQ |
|--------------------------|--|
| Purity | Greater than 95% as determined by reducing SDS-PAGE |
| Endotoxin level | <1 EU/µg as determined by LAL test. |
| Construction | Recombinant Mouse Adipocyte Complement-related 30 kDa Protein is produced by our E.coli expression system and the target gene encoding Glu18-Asn247 is expressed with a 6His tag at the N-terminus. |
| Accession # | Q60994 |
| Host | E.coli |
| Species | Mouse |
| Predicted Molecular Mass | 27.2 KDa |
| Formulation | Lyophilized from a 0.2 μ m filtered solution of PBS, pH 7.4. |
| Shipping | The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below. |
| Stability&Storage | Lyophilized protein should be stored at \leq -20°C, stable for one year after receipt. Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of reconstituted samples are stable at \leq -20°C for 3 months. |
| Reconstitution | Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. |

SDS-PAGE image



Background



Alternative Names

Background

Adiponectin; 30 kDa Adipocyte Complement-Related Protein; Adipocyte complement-related 30 kDa protein; ACRP30; Adipocyte; C1g and Collagen Domain-Containing Protein; Adipose Most Abundant Gene Transcript 1 Protein; apM-1; Gelatin-Binding Protein; ADIPOQ Adiponectin is a secreted protein. It is synthesized exclusively by adipocytes and secreted into plasma. Adiponectin is an important adipokine that is involved in the control of fat metabolism and insulin sensitivity, with direct anti-diabetic, antiatherogenic and anti-inflammatory activities. Adiponectin Stimulates AMPK phosphorylation and activates in the liver and the skeletal muscle, enhancing glucose utilization and fatty-acid combustion. Adiponectin also antagonizes TNFalpha by negatively regulating its expression in various tissues such as liver and macrophages, and also by counteracting its effects. It inhibits endothelial NFkappa-B signaling through a cAMP-dependent pathway. Adiponectin may play a role in cell growth, angiogenesis and tissue remodeling by binding and sequestering various growth factors with distinct binding affinities, depending on the type of complex: LMW, MMW or HMW.

Note For Research Use Only , Not for Diagnostic Use.