Product Name: Recombinant Mouse IFN alpha2

Catalog #: PEM0967



Summary

Name Interferon α-2/IFN α-2/IFN alpha2

Purity Greater than 95% as determined by reducing SDS-PAGE

Endotoxin level <1 EU/μg as determined by LAL test.

Construction Recombinant Mouse Interferon Alpha-2 is produced by our E.coli expression

system and the target gene encoding Cys24-Glu190 is expressed.

Accession # P01573

Host E.coli
Species Mouse

Predicted Molecular Mass 19.5 KDa

Formulation Lyophilized from a 0.2 µm filtered solution of 20mM Histidine-HCl, 6% Sucrose,

4% Mannitol, 0.02% Tween80 (w/v), pH 6.0.

Shipping The product is shipped at ambient temperature. Upon receipt, store it

immediately at the temperature listed below.

Stability&Storage Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt.


Reconstituted protein solution can be stored at 2-8°C for 2-7 days. Aliquots of

reconstituted samples are stable at \leq -20°C for 3 months.

Reconstitution Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is

not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles. Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image



Background

Product Name: Recombinant Mouse IFN alpha2



Catalog #: PEM0967

Alternative Names Interferon Alpha-2; IFN-Alpha-2; Interferon Alpha-A; LeIF A; IFNA2

Background At least 23 different variants of Interferon- α are known. The individual proteins

have molecular masses between 19-26 kD and consist of proteins with lengths of 156-166 and 172 amino acids. All IFN- α subtypes possess a common conserved sequence region between amino acid positions 115-151 while the amino-terminal ends are variable. Many IFN- α subtypes differ in their sequences at only one or two positions. Naturally occurring variants also include proteins truncated by 10 amino

acids at the carboxyl-terminal end.

Note

For Research Use Only, Not for Diagnostic Use.

Web: https://www.enkilife.com E-mail: order@enkilife.com techsupport@enkilife.com Tel: 0086-27-87002838