

Product Name: Recombinant Rat HER2 (C-6His)
Catalog #: PER0337




Summary

Name	HER2/CD340/ERBB2/Receptor Tyrosine-Protein Kinase ErbB-2
Purity	Greater than 95% as determined by reducing SDS-PAGE
Endotoxin level	<1 EU/μg as determined by LAL test.
Construction	Recombinant Rat Receptor Tyrosine-protein Kinase ErbB-2 is produced by our E.coli expression system and the target gene encoding Ala67-Val323 is expressed with a 6His tag at the C-terminus.
Accession #	P06494
Host	E.coli
Species	Rat
Predicted Molecular Mass	29.3 KDa
Formulation	Lyophilized from a 0.2 μm filtered solution of PBS, 5% Trehalose, 4M Urea, pH 7.4.
Shipping	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
Stability&Storage	Store at ≤-70°C, stable for 6 months after receipt. Store at ≤-70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.
Reconstitution	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100μg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.

SDS-PAGE image

Product Name: Recombinant Rat HER2 (C-6His)
Catalog #: PER0337



Alternative Names

Receptor tyrosine-protein kinase erbB-2; Epidermal growth factor receptor-related protein; Proto-oncogene Neu; Proto-oncogene c-ErbB-2; p185erbB2; p185neu; CD340; ERBB2

Background

ERBB2 belongs to the protein kinase superfamily, Tyr protein kinase family and EGF receptor subfamily. It contains a protein kinase domain. ERBB2 is widely expressed in epithelial cells, and amplification and/or overexpression of ErbB2 has been reported associated with malignancy and a poor prognosis in numerous carcinomas, including breast, prostate and ovarian cancers. Rat ERBB2 is an essential component of a neuregulin-receptor complex, although neuregulins do not interact with it alone. ErbB2 mediates signalling pathways which involve mitogen-activated protein kinase and phosphatidylinositol-3 kinase, this receptor plays a key role in development, cell proliferation and differentiation.

Note

For Research Use Only , Not for Diagnostic Use.